

Non Responsive Celiac Disease

Key Points

- Currently, the only treatment for celiac disease (CD) is strict life-long adherence to a gluten-free diet (GFD). It is not uncommon, however, for patients to fail to completely improve or relapse while on the diet.¹
- Non responsive Celiac Disease (NRCD) affects 20-30% of patients on a GFD.^{2,3} It may be defined as continued signs and symptoms that suggest CD despite dietary gluten avoidance for 6-12 months.⁴
- NRCD can be further classified as primary or secondary. Primary refers to an initial failure to achieve symptomatic response on a GFD. Secondary NRCD occurs when symptoms return while on a GFD.²
- When the symptoms fail to improve or recur, proper evaluation should be followed to identify and treat the specific cause (See Figure 1).⁵ It is important to review and carefully examine the initial diagnosis of CD based on the history of symptoms, signs, lab tests, and biopsy results at the time of diagnosis.
- An endoscopy with small intestinal biopsy may be considered depending on symptoms and/or blood test results. This will help the doctor assess intestinal healing and other conditions that can cause similar biopsy findings. Biopsies of the small intestine are obtained with the help of a safe and quick procedure called an esophago-gastro-duodenoscopy (EGD) or endoscopy. (See [Blood Tests and Endoscopy](#)).
- The most common cause of NRCD is accidental gluten ingestion.³ Gluten exposure accounts for 35-50% of ongoing symptoms in patients with CD.⁴ For this reason, it is very important to obtain sound dietary advice from an expert celiac dietitian.¹ It will also help you identify if the cause of ongoing symptoms could be due to lactose intolerance or fructose malabsorption. These two conditions can occur as a result of intestinal damage caused by CD.
- A positive celiac serology blood test for CD (IgA-tTG) could be helpful to assess cross-contact with gluten and adherence to the GFD. It is important to remember, however, that a normal celiac serology does not exclude cross contact with gluten as a possibility.
- After a dietitian carefully excludes gluten as a cause of the symptoms, a small intestinal biopsy may be repeated and compared with the initial biopsy. A normal or near normal biopsy may lead the doctor to consider other possible causes (See Table 1).⁶

Other causes of NRCD besides ongoing gluten exposure include:

1. Small intestinal bacterial overgrowth syndrome (SIBO):

- Small intestinal bacterial overgrowth syndrome (SIBO) is caused by abnormal intestinal motion, lowered immune defenses, or damage to the intestinal mucosa. These factors create imbalance in the gut flora that results in overgrowth of the harmful bacteria.
- This condition can be challenging but is often assessed with a hydrogen and methane breath test and treated with a course of antibiotics. The use of prebiotics and probiotics may also be useful.

2. Microscopic colitis:

- Microscopic colitis is a common cause of long-standing diarrhea caused by inflammation of the large intestine (colon).
- It generally presents with episodes of watery, non-bloody diarrhea of long standing, intermittent, or recurrent course.⁷
- The diagnosis is made by colonoscopy and confirmed by assessment of biopsies under a microscope (thus the name microscopic).
- Treatment varies depending on the persistence and severity of symptoms.

3. Pancreatic insufficiency:

- Pancreatic insufficiency is caused by insufficient production of digestive enzymes by the pancreas. This low production leads to malabsorption of nutrients and fatty diarrhea (pale, bulky and foul smelling stools, often with oil droplets).
- Diagnosis is made by measuring the amount of enzymes (usually elastase) and/or fat in the stool.
- Treatment is based primarily on pancreatic enzyme replacement therapy. It may also include lifestyle modifications and vitamin supplementation.

4. Irritable bowel syndrome (IBS):

- IBS is a functional disorder of the intestine that commonly presents with abdominal pain or discomfort, is associated with bowel changes and is usually relieved by bowel movements. Other symptoms such as bloating, diarrhea and/or constipation are required to establish a diagnosis.
- No damage is seen in the intestine compared to CD.
- In many cases, this condition can be controlled by changes in lifestyle including diet and stress management.

5. Refractory Celiac Disease (RCD):^{4,8,9}

- RCD is a severe form of CD that belongs to a subset of NRCD.
- This is a rare condition affecting 1-2% of patients with CD and up to 10% of those with NRCD. Damage to the small intestine continues despite a strict GFD and no evidence of another disease, including cancer (lymphoma).

- RCD can be further subdivided into type I and type II. This distinction is based on a special type of white blood cells seen on the biopsy of the small intestine.
- RCD type I has a better clinical course and survival than RCD type II.

Other diagnoses that may be associated with CD include inflammatory bowel disease and lactose or fructose intolerance (caused by mucosal damage from CD). These conditions should be excluded and treated correctly.¹

Take Home Messages:

- In the majority of people with CD, a GFD is sufficient to allow for clinical improvement and improved biopsy results.
- Some patients however will still have symptoms 6-12 months after starting a GFD or have recurrent symptoms after improving initially on a GFD.
- The most common cause of ongoing symptoms and abnormal lab values is cross contact with gluten.
- GFD adherence should be supported by the guidance of an expert dietitian, advocacy groups, and regular clinic visits.
- Treatment of other causes of NRCD depends on the cause.

Table 1:


Causes of Nonresponsive Celiac Disease

Cause	Typical Features/Tests	How Common
Gluten exposure (see Chapter 20)	Evaluation by dietitian skilled in celiac disease	Very common
Irritable bowel syndrome (see Chapter 38)	None	Very common
Lactose intolerance or fructose malabsorption (see Chapter 37)	Trial of lactose or fructose restriction; lactose or fructose breath testing	Somewhat common
Microscopic colitis (see Chapter 39)	Biopsy of colon	Somewhat common
Small intestinal bacterial overgrowth (see Chapter 40)	Breath testing and or a response to antibiotic therapy	Somewhat common
Refractory celiac disease (see Chapter 43)	Biopsy of small intestine	Rare
Eating disorder (see Chapter 30)	None	Rare
Inflammatory bowel disease	Biopsy of small or large intestine, imaging studies of intestine	Rare
Pancreatic exocrine insufficiency	Stool levels of chymotrypsin or elastase	Rare
Motility disturbances (too slow or too fast movement of food through the intestine)	Gastric emptying study, intestinal transit testing	Rare
Food allergy (see Chapter 35)	Allergy testing (skin or blood)	Very rare
Cancer	Endoscopy, imaging studies of intestine	Very rare

From Shailaja Jamma, MD, and Daniel A. Leffler, MD: Nonresponsive Celiac Disease. In *Real Life with Celiac Disease: Troubleshooting and Thriving Gluten Free* by Melinda Dennis, MS, RD, LDN, and Daniel A. Leffler, MD.


www.reallifewithceliacdisease.com

Figure 1: An approach to the investigation of Nonresponsive celiac disease



Supporting evidence

- Confirmation of small bowel histology findings consistent with celiac disease
- Positive TTG, DGP or EMA serology at some time during the clinical course
- Presence of HLA DQ2 or DQ8
- Biopsy proven dermatitis herpetiformis
- Clinical and/or histological response to GFD
- Family history of CD



References:

1. Baggus EMR, Hadjivassiliou M, Cross S, Penny H, Urwin H, et al. How to manage adult coeliac disease: perspective from the NHS England Rare Diseases Collaborative Network for Non-Responsive and Refractory Coeliac Disease. *Frontline Gastroenterol.* 2019 Aug 8;11(3):235-242.
2. Leffler D, Dennis M, Hyett B, Kelly E, Schuppan D, Kelly C. Etiologies and predictors of diagnosis in nonresponsive celiac disease. *Clin Gastroenterol Hepatol.* 2007; 5(4): 445-450.
3. Aggarwal N, Bhatia U, Dwarakanathan V, Singh AD, Singh P, Ahuja V, Makharia GK. Prevalence and etiologies of non-responsive celiac disease: A systematic review and meta-analysis. *J Gastroenterol Hepatol.* 2025 Jan;40(1):101-107.
4. Rubio-Tapia, A, Hill ID, Kelly CP, Calderwood AH, Murray JA; ACG clinical guidelines: diagnosis and management of celiac disease. *Am J Gastroenterol.* 2013;108(5):656-676.
5. Abdulkarim, A, Abdulkarim AS, Burgart LJ, See J, Murray JA. Etiology of nonresponsive celiac disease: results of a systematic approach. 2002. *Am J Gastroenterol* 97(8):2016-2021.
6. Dennis, M, Leffler D. In *Real Life with Celiac Disease: Troubleshooting and Thriving Gluten Free.* AGA Press. Bethesda, MD, 2010.

7. Storr MA. Microscopic colitis: epidemiology, pathophysiology, diagnosis and current management-an update 2013. *ISRN Gastroenterol* 2013; 352718.
8. Hollon JR, Cureton PA, Martin ML, Puppa EL, Fasano A. Trace gluten contamination may play a role in mucosal and clinical recovery in a subgroup of diet-adherent non-responsive celiac disease patients. 2013. *BMC Gastroenterol*;13:40.
9. Dewar, DH, Donnelly SC, McLaughlin SD, Johnson MW, Ellis HJ, Ciclitira PJ. Celiac disease: management of persistent symptoms in patients on a gluten-free diet. *World J Gastroenterol*. 2012;18(12):1348-1356.
